


## Review Activity Module 6: Ecology

Laroche:


### Disappearing Sea Otters

Dr. James Estes has studied sea otters off the coast of Alaska since the 1970s. Once an endangered species as a result of fur trading, otter populations recovered and appeared very healthy and robust in the early 1980s. However, starting in the early 1990s, Estes and his colleagues noted that otter population numbers were dropping precipitously.



Changes in relative abundance of sea otters at several locations in the Aleutian Islands, Alaska.

(Redrawn from Estes et al., 1998.)



Map of the North Pacific Ocean showing the Aleutian Islands and some specific sea otter study sites. (From Estes, J.A., and D.O. Duggins. 1995)

The following excerpt from the New York Times describes the thought process and background research done by Estes and colleagues to examine this question:

*Could the otters simply have migrated from one part of the region to another? To find out, the researchers analyzed populations over a 500-mile-long stretch of the Aleutians from Kiska to Seguam.... By 1993 otter numbers in that whole stretch had been cut by half. Here the geographical scope of the research effort became critical; a smaller region would not have been large enough to reveal the decline. In 1997, they ... found that the population decline had worsened, to about 90 percent....*

*"That told us for sure it was a very large- scale decline, but we were still trying to understand the cause," Dr. Estes said.... The researchers ... ruled out reproductive failure. Their studies enabled them to keep track of how often otters gave birth and how many young survived, and this revealed that reproduction was continuing to re-supply the population.*

*With other possible causes eliminated, ... mortality had to be the explanation. In the past, they had seen temporary declines in otter populations because of starvation, pollution or infectious disease. "In all those cases," Dr. Estes said, "we find lots of bodies. They get weak and tired and come ashore to die." This time not a single dead otter was found -- a clue, he said, that "something really weird was going on."*

(Excerpted from Stevens, William K. "Search for missing sea otters turns up a few surprises." *New York Times*, January 5, 1999.)

1. Based on the information presented above, propose a hypothesis for what could be causing the decline in sea otter numbers.


**If mortality has to be causing the decrease, but no bodies are being found, then it stands to reason that something is consuming these bodies, or, rather, killing and eating the otters. The only organisms that could and would do this in Alaska are killer whales or Greenland sharks.**




<https://www.youtube.com/watch?v=k5Eo2FGeEyg>

Estes and his group hypothesized that increased predation by killer whales was the cause of the sea otter decline. This was an unusual idea, since killer whales and sea otters had been observed together in Alaska for decades with no obvious interactions occurring between them. The first time a killer whale was observed attacking a sea otter was in 1991. Nine more attacks were observed in the next seven years and it was these attacks that finally led Dr. Estes and his colleagues to propose their hypothesis.

They estimated the impact of killer whales on sea otter populations by comparing trends in population size and survival rates of individually marked otters between two adjacent locations on Adak Island: Clam Lagoon and Kuluk Bay. Kuluk Bay is on an open coast, so sea otters there are exposed to killer whales. In Clam Lagoon, the entrance from the open sea is too narrow and shallow for killer whales to get in.



Changes in sea otter population size over time at Clam Lagoon and Kuluk Bay, Adak Island, Alaska. Redrawn from Estes et al., 1998.



Survival rates of sea otters individually marked in 1995 with flipper tags and radio transmitters at Clam Lagoon and Kuluk Bay, Adak Island, Alaska. Redrawn from Estes et al., 1998.

2. Based on the two figures above, what can you conclude about the effects of killer whales on sea otter populations?

**The colony subjected to killer whales displays much greater mortality and a sharp population decline, while the “control” population not subjected to killer whales remains constant, with far less mortality. This clearly suggests that killer whales are having a substantially negative effect on otter population numbers.**

#### **Killer whale and sea otter energetics**

Estimated number of Aleutian Island sea otters eaten, 1990-1996	40 000
<b>Adult sea otters</b>	
Average caloric content	1.81 kcal/gram wet weight
Average mass, male	34 kg
Average mass, female	23 kg
<b>Killer whales</b>	
Average field metabolic rate	55 kcal/kg of whale/day
Average mass, male	5600 kg
Average mass, female	3400 kg

Using the table above, it's possible to calculate that only 3.6 male killer whales, feeding exclusively on otters, would be able to account for the decrease in otter population over the six years (try to calculate this number yourself). Of course, killer whales would not feed exclusively on otters, but, conversely, there are much more than just 4 killer whales patrolling the Aleutian Islands. Thus, it seems entirely reasonable that killer whales could account for the decrease in otter numbers.

This, of course, begs the very important question: Killer whales and sea otters had co-existed peacefully for decades prior to the 1990s, so why would the whales suddenly start eating otters?

3. Propose at least one, and maybe several different hypotheses for why whales would begin eating otters.

**If whales are ‘suddenly’ eating otters, it’s because these have become the best food option for them to choose. It’s unlikely that otters somehow just became more palatable; what’s much more likely is that another preferable food source is no longer available, and so killer whales have to switch to whatever they can get, or, rather, whatever they encounter. Since otter populations were initially high, it’s likely that whales encountered these fairly frequently. However, with other preferable prey around, the whales simply ignored the otters. Now, that’s no longer the case.**

Estes et al. (1998) proposed a hypothesis to address this question. Their hypothesis incorporated the following elements:

- The species composition of Bering Sea fish changed dramatically in the decades preceding the 1990s. Oily fish species, such as ocean perch (a rockfish) and herring, declined sharply, while the abundance of pollock, a less oily fish, increased. The causes of these changes in the fish community are not clear. Several plausible hypotheses have been proposed, including overexploitation of perch and herring by humans through the 1960s and the warming of the northern Pacific in the 1970s, which may have made the temperature unsuitable for some species. Human hunting of plankton-feeding baleen whales may even indirectly play a role. Since pollock are also plankton feeders, a reduction in whales may have allowed pollock to proliferate.
- Steller sea lion and harbor seal populations in the North Pacific collapsed starting in the late 1970s. The decline in oily fish species may have contributed to this collapse, because, although pollock were increasingly available as an alternate food and are eaten by sea lions and seals, they are a less nutritious food and may not provide sufficient nutrition, especially for the growing juveniles. Whatever the cause, North Pacific harbor seal and sea lion populations had declined by as much as 80 percent by 1992.
- Killer whales are found in all oceans and seas from the polar regions to the tropics. They live in groups called pods, which appear to be of two genetically distinct types: transients and residents. Transients form smaller pods (one to seven individuals), roam over broader areas, eat predominately mammals (including seals, sea lions, and otters, but also other porpoises, and even large baleen whales, which they hunt and attack as a group), and vocalize (using a series of whistling noises) less. Residents form pods of five to 25 individuals, have smaller ranges, eat mostly fish, and vocalize frequently.

- Seals and sea lions (together called pinnipeds) insulate themselves in the cold ocean using a thick layer of blubber (fat), while otters insulate themselves based on two layers of hair, one of which traps air. As a result, pinnipeds have an average caloric content of 5.35 kcal/gram wet weight, as compared to 1.81 kcal/gram wet weight for otters. The pinnipeds are also much larger than the otters, with steller sea lion females averaging 300 kg (still MUCH smaller than a 4000 kg killer whale)
4. Using this information, apply optimal foraging theory and a cost-benefit approach to analyzing animal behaviour to either refine one of your hypotheses from above or propose an entirely new hypothesis for why killer whales may have begun eating otters in the 1990s. In your response, **compare and contrast the basic costs and benefits for an individual killer whale when choosing to eat an otter as compared to other mammal (or pinniped) prey**, and then hypothesize why whales would make this choice.

**Given the still massive size discrepancy between a killer whale and a seal or sea lion, it's unlikely that catching or killing these animals is any more difficult than catching or killing an otter. However, for the effort required to kill a sea lion, there is substantially more reward than eating an otter, both because the sea lion is larger, but also because the animal has a thick and energy rich layer of fat. Thus, it would seem optimal for transient killer whales to hunt pinnipeds preferably over otters. When populations of pinnipeds crashed, however, the result was a scenario of low prey abundance. Under these conditions, predators are expected to eat whatever they come across in order to meet their metabolic needs. Since otter populations were healthy, hungry whales likely encountered these frequently, and were forced to eat them in the absence of their more rewarding pinniped prey.**

5. Based on your hypothesis and the cost benefit analysis above, make a prediction for how killer whale population numbers might have changed in the late 1990s and early 2000s. Explain the rationale behind your prediction.

**Transient killer whale populations likely decreased as well, because the cost-benefit ratio of eating otters is less favourable, resulting in an overall worse energy trade-off for individual whales. With less energy, there is less reproduction and more mortality, which leads to decreases in population size.**

In the early 2000s, and after over a decade of concerted conservation efforts, steller sea lion populations recovered somewhat, and stabilized at about 50% of pre-1970s levels. Based on this recovery, and with otter population levels so low, it seems that killer whales stopped hunting otters, or, at the very least, dramatically reduced their predation pressure. Consequently, otter populations also began to recover.


Below are demographic metrics for the Kuluk Bay sea otter population. In 2001, the population consisted of 36 individuals. For the sake of this exercise, we will ignore E and I for population growth calculations.

Year	B	D
2001	15	4
2002	20	5
2003	25	8
2004	29	11

6. Based on the data in the table above, complete the following table:

Year	$N$	$r$
2001	36	<b>0.31</b>
2002	<b>47</b>	<b>0.32</b>
2003	<b>62</b>	<b>0.27</b>
2004	<b>79</b>	<b>0.22</b>
2005	<b>97</b>	

7. Use the data from the table you just completed to plot a graph of otter population numbers in Kuluk Bay over the five year time period. Place this graph in the space below.



8. What kind of population growth (approximately) is being demonstrated in the graph above? Why do you believe the population grew this way during the five year span?

**The growth is initially exponential, although by years 3 and 4 it seems to be decreasing somewhat. Still, over the four years the population grew very well. This occurred because the limiting factor on population size, i.e., whale predation, was removed. Without whale predation keeping population numbers low, the otters had abundant resources and space, which made it possible to survive and reproduce at a high rate during the measured period of time.**

9. Would the population continue growing this way indefinitely? If yes, what would allow it to do so? If not, what would prevent it from doing so?

**No, the population could not grow exponentially indefinitely. Eventually the population would hit a carrying capacity, where other density limits like food and space would reduce overall birth rates and increase death rates to the point where these balanced out, resulting in zero growth and a reasonably stable population size.**

10. Assuming no other changes occur in the populations of other species (whales, sea lions, oily fish, etc.) from 2000 levels, and given all of the information presented above, would you predict that the otter population in Kuluk Bay and those in other areas would eventually grow to reach pre-1990s numbers, i.e., population numbers before whales began preying on otters? Explain your answer.


**I would predict that otter numbers would not increase to pre-predation levels. I would make this prediction because I don't think killer whales would stop feeding on otters entirely. While these would still remain a suboptimal food source, the pinniped populations have not recovered enough to fully support the killer whale populations, which means that these individuals would still need to eat otters on occasion. By imposing a certain amount of predation pressure on the otters, killer whales would likely reduce the realized carrying capacity of the environment down from where it was pre-1990, which means that otter numbers would likely not get as high.**

The following five pages present information about and display data from a long-term and large-scale study of sea otters and kelp forest communities in southeast Alaska and the Aleutian Islands.




Kelps are large brown algae (protists), and a kelp forest is a large stand of individual kelps that grow to the surface and form a floating canopy. Kelp forests are highly productive ecosystems: productivity can be as high as 1500-3000 g Carbon/m<sup>2</sup>/year, which is comparable to the most productive terrestrial and aquatic ecosystems. Also, the tall (up to 30 m) bodies of the kelp individuals create a complex three-dimensional structure that supports a very diverse community of associated algal and animal species. For example, California kelp forests are home to up to 750 kinds of fish and invertebrates, including a number of commercially and recreationally harvested species.

Sea urchins are also mentioned in these data. Urchins are small, spiny, globular animals which are members of the echinoderm phylum (like sea stars). They thus have many of the same characteristic echinoderm features, like pentaradial symmetry (obvious from underneath; not so much from above or in the picture) and a water-vascular system. They have sharp and pointy spines, which can inflict painful wounds if they penetrate human skin (I know from firsthand experience). Urchins feed primarily on algae, and are voracious kelp-munchers.



The graphs below and tables on the following **two** pages compare sea urchin and kelp abundances in areas with and without sea otters (see the map on the first page for locations). Although sea otters formerly were found at all of these locations, they were exterminated from most of their range by hunting during the 19<sup>th</sup> century. Amchitka and Adak Islands in the Aleutians were locations of some of the few remnant populations at the time otters were protected in the early 1900s. Sea otters were then re-introduced to southeast Alaska in 1968-71. This new population expanded into Surge Bay by the early 1970s and into Torch Bay in 1985.



Kelp density (individuals/0.25 m<sup>2</sup>) plotted against estimated sea urchin biomass (g/0.25 m<sup>2</sup>) for the Aleutian Islands and southeast Alaska. Points represent averages for sites at each location. Sea urchin biomass was estimated from samples of population density, size-frequency distribution, and the relationship between urchin diameter and wet mass. (From Estes, J.A., and D.O. Duggins. 1995)

Abundance and population characteristics of kelps and sea urchins at two locations in the Aleutians, Amchitka and Shemya Islands, in 1972 and 1987 (shown as means  $\pm$  1 standard error). The same four sites at Amchitka and two sites at Shemya were sampled in both years\*. Sea otters were continuously abundant at Amchitka and absent from Shemya during the 15-yr period. (From Estes, J.A., and D.O. Duggins. 1995.)

	Amchitka Island		Shemya Island	
	1972	1987	1972	1987
<b>Kelp species (inds./0.25 m<sup>2</sup>)</b>				
<i>Alaria fistulosa</i>	1.6 $\pm$ 1.30	0.3 $\pm$ 0.22	0	0.5
<i>Laminaria</i> spp.	2.3 $\pm$ 0.49	3.9 $\pm$ 0.95	0	0
<i>Agarum cribrosum</i>	1.2 $\pm$ 0.61	0.5 $\pm$ 0.42	0	0
<i>Thalassiophyllum clathrus</i>	0.1	0	0	0
<b>Total kelps</b>	5.1 $\pm$ 0.66	4.7 $\pm$ 1.15	0	0.5
<b>Sea urchins</b>				
Maximum test diameter (mm)	30.5 $\pm$ 1.34	27.3 $\pm$ 3.24	72.5 $\pm$ 0.71	70.5 $\pm$ 4.95
Biomass (g/0.25 m <sup>2</sup> )	45.1 $\pm$ 16.9	36.7 $\pm$ 15.0	368.2 $\pm$ 151.7	369.3 $\pm$ 14.3
Density (inds./0.25 m <sup>2</sup> )	27.9 $\pm$ 14.5	23.4 $\pm$ 7.5	50.0 $\pm$ 14.6	38.6 $\pm$ 1.4

\*The 1972 data were obtained from 10 haphazardly placed 0.25-m<sup>2</sup> quadrats/site, the 1987 data from 20 randomly placed 0.25-m<sup>2</sup> quadrats/site.

Abundance and population characteristics of kelp and sea urchins at two locations in southeast Alaska, Torch Bay (1976-1978) and Surge Bay (1978 and 1988), shown as means  $\pm$  1 standard error. Sea otters were continuously absent at Torch Bay and present at Surge Bay during these time periods. (From Estes, J.A., and D.O. Duggins. 1995.)

	Torch Bay			Surge Bay	
	1976	1977	1978	1978	1988
<b>Kelps (inds./m<sup>2</sup>)</b>					
Annuals*	2.1 $\pm$ 1.39	0.2 $\pm$ 0.25	11.6 $\pm$ 6.69	2.1 $\pm$ 0.45	3.7 $\pm$ 2.34
Perennials**	0.1 $\pm$ 0.11	0	0.9 $\pm$ 1.14	48.4 $\pm$ 6.33	50.3 $\pm$ 7.46
<b>Total</b>	2.2	0.2	12.5 $\pm$ 5.56	50.5 $\pm$ 6.43	54.0 $\pm$ 9.33
<b>Sea urchins (inds./m<sup>2</sup>)</b>					
<i>S. franciscanus</i>	3.6 $\pm$ 3.05	3.8 $\pm$ 2.55	4.9 $\pm$ 3.71	0	0
<i>S. purpuratus</i>	1.0 $\pm$ 0.75	2.3 $\pm$ 2.52	0.3 $\pm$ 0.41	0	0
<i>S. droebachensis</i>	3.4 $\pm$ 2.24	1.5 $\pm$ 0.95	0.2 $\pm$ 0.18	0.02	0.04
<b>Total</b>	8.0 $\pm$ 4.56	7.6 $\pm$ 5.78	5.4 $\pm$ 4.27	0.02	0.04

\*Primarily *Alaria fistulosa* and *Nereocystis leutkeana*.

\*\*Primarily *Laminaria groenlandica*.

The table below summarizes data from several studies on the diet of sea otters.

Occurrence of prey items in sea otter stomachs and feces. (From Estes, J.A., N.S. Smith, and J.F. Palmisano. 1978. "Sea otter predation and community organization in the western Aleutian Islands, Alaska." *Ecology* 59:822-833.)

<b>Source</b>	<b>Wilke 1957</b>	<b>Kenyon 1969</b>	<b>Kenyon 1969</b>	<b>Burgner and Nakatani 1972</b>	<b>Barahash- Nikiforov 1947</b>	<b>Williams 1938</b>
<b>Location</b>	Amchitka	Amchitka	Amchitka	Amchitka	Commander Islands	Western Aleutians
<b>Sample period</b>	1954	1962-1963	1962-1963	1970	1930-1932	1936
<b>Sample type</b>	Stomach	Stomach	Stomach	Stomach	Feces	Feces
<b>Sample size</b>	5	309	309	49	500	70
<b>Analysis</b>	<b>Percent of total volume</b>	<b>Percent of total volume</b>	<b>Percent of total number of prey item</b>	<b>Percent of stomachs containing food item*</b>	<b>Percent of total volume</b>	<b>Percent of total volume</b>
<b>Prey item</b>						
Annelids	0	1	2	2	0	0
Arthropods						
Crabs	0	< 1	4	22	10	4
Others	0	0	3	0	0	0
Mollusks	8	37	31	38	23	13
Echinoderms						
Sea urchins	86	11	21	82	59	78
Others	0	0	16	0	0	0
Fish	6	50	22	44	7	3
Others	0	< 1	1	0	1	2
<b>Total</b>	100	100	100	-	100	100

\*Percent of total volume: carnivores 65 (including fish 62.2) and herbivores 35.

11. What do these data tell you about the role of sea otters in their community? Include in your answer a description of how you think sea otters are affecting the two other groups of species (urchins and kelp), as well as a description of what effect you expect sea otters to have on the rest of the kelp forest community.


**Sea otters are keystone species in their communities. This means that their overall effect on community composition is out of proportion with their abundance in their community. Their effect on the community is to eat sea urchins. Without otters, sea urchin populations explode, and the kelp forests are decimated, in that urchins eat the kelp. By keeping urchin numbers low, otters make the kelp forests possible, which in turn provides an environment that supports myriad other species. So, without otters, urchins eat the kelp, the forest disappears, and a wide variety of fish, algal, and invertebrate species go as well. This is why otters are a keystone species.**

The table below displays all of the species or groups of species encountered so far in this ‘story’, along with their approximate mass per individual, and estimated numbers of individuals found within the same 10 Km<sup>2</sup> area surrounding Kuluk Bay. Phytoplankton consists of photosynthetic bacteria and microscopic protists; “kelp fish” is a broad category, and will be considered as fish that eat kelp or other algae, while being eaten by bigger fish like perch or pollock. Several of the animal species are transient, so estimates are of the number of given organisms of this species within the area at any given time. Use the data from this table to answer the next two questions.


<b>Organism</b>	<b>Mass per Individual (Kg)</b>	<b>Number Estimated</b>
Killer Whales	4500	4
Sea Lions	400	5
Seals	200	10
Otters	28.5	200
Ocean Perch	1.5	1500
Herring	3	2000
Pollock	2	7500
Kelp Fish	0.5	15000
Urchins	0.1	1000
Kelp Invertebrates	0.000001	1E+11
Baleen Whales	36000	2
Kelp	5	15000
Phytoplankton	0.000000001	1E+15

12. Draw a food web of the broad Aleutian Islands aquatic community. Include in the diagram all of the species or groups of species encountered so far: otters, killer whales, ocean perch, herring, Pollock, steller sea lions, harbor seals, baleen whales, plankton (let's say just the photosynthetic phytoplankton), kelp, kelp fish (yes, this is extremely broad; let's just say the "kelp fish" eat kelp or other algae, and are eaten by bigger fish like perch or pollock), kelp invertebrates, and urchins. In your web, identify each organism as a primary producer (pp), a primary consumer (1), a secondary consumer (2), etc.

**I don't have the time to actually draw a web, so take a look at this:**



13. Using the data from the table above, draw a biomass pyramid for the aquatic ecosystem. In your pyramid, be sure to indicate the number of organisms you've assigned to each level, the overall mass of organisms at each level, and the approximate energy transfer efficiency between each level.



The energy transfer rates calculated for the pyramid below do not directly reflect a straight transfer between levels. For example, all of the biomass contained in baleen whale primary consumers is not available for secondary consumers (and, frankly, some of these secondary consumers are actually eaten by baleen whales... that's not really important here). Also, the biomass of quarternary consumer whales is higher than the biomass of tertiary consumers, but is supported by the fact that the whales eat tertiary, secondary, and primary consumers (including the VERY large baleen whales). If you add the biomass of all of these organisms up, you get the 22% energy transfer rate.

In the end, calculating energy transfer rates is very tricky, and there are numerous answers that could conceivably be correct. Here, while it wouldn't be technically correct to simply divide the mass at a higher trophic level by the mass beneath it, it is an answer I would accept on a test.

You should also note that this pyramid exclusively reflects mass, not biomass, which is a measure of biological density. Since all of these mass measurements were taken for the same area, the density measure was simply ignored. This also isn't a productivity pyramid (g biological matter/ unit area / unit time), and so doesn't really properly describe energy transfers.


## Roffey:

### Ecology Exit Quiz

1. All of the following are density-dependent factors that limit animal populations EXCEPT (note: I'm pretty confident that this is the correct answer, but neither birthrate nor mortality are density-dependent *factors*; rather, they are characteristics of a population that can be affected by density-dependent factors)
  - A. weather
  - B. predation
  - C. birthrate
  - D. food competition
  - E. mortality
2. During the carbon cycle, which of the following carbon compounds would be utilized as an energy source by heterotrophs?
  - A. calcium carbonate
  - B. carbonic acid
  - C. organic molecules
  - D. carbon dioxide
  - E. carbon monoxide
3. All of the following statements concerning characteristics of predator-prey relationships are correct EXCEPT:
  - A. A rise in the population of prey is often followed by a rise in the population of predators.
  - B. A rise in the population of predators is followed by a decrease in the population of prey.
  - C. Camouflage is an adaptation that protects prey.
  - D. The production of large numbers of offspring within very short periods of time ensures the survival of some prey populations.
  - E. The population of predators most often eliminates the population of prey.
4. Which of the following is true about secondary consumers in an ecosystem?
  - A. They eat only plants.
  - B. They are eaten by primary consumers.
  - C. They are smaller and weaker than are primary consumers.
  - D. They are fewer in number than are primary consumers.
  - E. They contain the greatest total biomass in the system.
5. In the nitrogen cycle, the transformation of gaseous nitrogen into nitrogen-containing compounds is performed primarily by
  - A. fungi
  - B. bacteria
  - C. green plants
  - D. herbivores
  - E. carnivores

6. Which point on the curve in the diagram to the right best represents the carrying capacity of the environment for the population shown?

- A. A
- B. B
- C. C
- D. D
- E. E




7. Which of the following organisms is most likely to be located at the apex of the pyramid of biomass?

- A. grass
- B. grasshopper
- C. snake
- D. mouse
- E. hawk

8. All of the following statements about the diagram are correct EXCEPT:

- A. The grasshopper is an herbivore.
- B. Only two trophic levels are depicted.
- C. The mouse and grasshopper are at the same trophic level.
- D. The grass is a producer.
- E. All of the organisms except grass are consumers, regardless of position.



9. The organic and inorganic materials in all the organisms in the diagram will eventually return to the environment by the action of

- A. decomposers
- B. producers
- C. primary consumers
- D. secondary consumers
- E. top carnivores

10. Which of the following best explains why there are seldom more than five trophic levels in a food chain?

- A. Most carnivores function at more than one trophic level.
- B. Trophic levels above this number contain too many individuals.
- C. Top carnivores are too few in number to prey effectively.
- D. The ecosystem contains too much biomass.
- E. Energy is lost from each trophic level.